

Complex Analysis

Chao Ren

January 31, 2026

Contents

1	Vector Spaces and Linear Maps	2
1.1	Vector Spaces	2
1.2	Linear Maps	3

Chapter 1

Vector Spaces and Linear Maps

1.1 Vector Spaces

Definition 1. A set \mathbb{F} with two binary operations $+$ and \times is a **field** if both $(\mathbb{F}, +, 0)$ and $(\mathbb{F} \setminus \{0\}, \times, 1)$ are abelian groups and the distribution law holds:

$$(a+b)c = ac + bc, \text{ for all } a, b, c, \in \mathbb{F}$$

Definition 2. The smallest integer p such that

$$1 + 1 + \cdots + 1 \quad (p \text{ times}) = 0$$

is called the **characteristic** of \mathbb{F} . If no such p exists, the characteristic of \mathbb{F} is defined to be 0. If such a p exists, it is necessarily prime.

Definition 3. A **vector space** V over a field \mathbb{F} is an abelian group $(V, +, 0)$ together with a scalar multiplication $\times \mathbb{F} \times V \rightarrow V$ such that for all $a, b \in \mathbb{F}, v, w \in V$:

1. $a(v + w) = av + aw$
2. $(a + b)v = av + bv$
3. $(ab)v = a(bv)$
4. $1 \cdot v = v$

Definition 4. Let V be a vector space over \mathbb{F} .

1. A set $S \subseteq V$ is **linearly independent** if whenever $a_1, \dots, a_n \in \mathbb{F}$,

and $s_1, \dots, s_n \in S$,

$$a_1 s_1 + \dots + a_n s_n = 0 \Rightarrow a_1 = \dots = a_n = 0.$$

2. A set $S \subseteq V$ is **spanning** if for all $v \in V$ there exists $a_1, \dots, a_n \in \mathbb{F}$ and $s_1, \dots, s_n \in S$ with $v = a_1 s_1 + \dots + a_n s_n$.
3. A set $\mathcal{B} \subseteq V$ is a **basis** of V if \mathcal{B} is spanning and linearly independent. The size of \mathcal{B} is the **dimension** of V .

Example. Let $V = \mathbb{R}^{\mathbb{N}} = \{(a_1, a_2, a_3, \dots) \mid a_i \in \mathbb{R}\}$. Then $S = \{e_1, e_2, \dots\}$ where $e_1 = (1, 0, 0, \dots), \dots$ is linearly independent but its span W is a proper subset of V .

Proof. It is important that n is finite.

1.2 Linear Maps

We consider linear maps and their relation to matrices.

Definition 5. Suppose V and W are vector spaces over \mathbb{F} . A map $T : V \rightarrow W$ is a **linear map** if for all $a \in \mathbb{F}, v, v' \in V$,

$$T(av + v') = aT(v) + T(v').$$

A bijective linear map is called an **isomorphism** of vector spaces.